

**DETERMINATION OF BUNCH COMPONENT WHICH CONTRIBUTE
WITH OIL-TO-BUNCH (O/B) RATIO OF COMMERCIAL PRODUCTION
IN SIME DARBY BANTING**

MUHAMMAD SHUKRI BIN JUMALI

**Final Year Project Report Submitted in
Partial Fulfilment of the Requirements for the
Degree of Bachelor of Science (Hons.) Plantation Technology and Management
in the Faculty of Plantation and Agrotechnology
Universiti Teknologi MARA**


JULY 2015

DECLARATION

This Final Year Project is a partial fulfilment of the requirements for a degree of Bachelor of Science (Hons.) Plantation Technology and Management, Faculty of Plantation and Agrotechnology, Universiti Teknologi MARA.

It is entirely my own work and has not been submitted to any other University or higher education institution, or for any other academic award in this University. Where use has been made of the work of other people it has been fully acknowledged and fully referenced.


I hereby assign all and every right in the copyright to this Work to the Universiti Teknologi Mara ("UiTM"), which henceforth shall be the owner of copyright in this Work and that, any reproduction or use in any form or by any means whatsoever is prohibited without a written consent of UiTM.

Candidate's signature: 

Date: 10/08/2015

Name: MUHAMMAD SHUKRI BIN JUMALI

I hereby declare that I have checked this project and in my opinion, this project is adequate in terms of scope and quality for the award of the degree of Bachelor of Science (Hons.) Plantation Technology and Management, Faculty of Plantation and Agrotechnology, university Teknologi MARA.


Signature:

Name: Prof. Dr. MOHAMAD OSMAN

Position: LECTURER

Date: 10/08/2015

TABLE OF CONTENTS

	<u>Page</u>
ACKNOWLEDGEMENT	Iii
TABLE OF CONTENT	iv
LIST OF FIGURES	v
LIST OF TABLES	vi
LIST OF ABBREVIATIONS	vii
ABSTRACT	viii
ABSTRAK	ix
 <u>CHAPTER</u>	
1 INTRODUCTION	
1.1. Background	1
1.2. Problem statement	2
1.3. Research question	2
1.4. Objective of study	3
1.5. Significant of study	3
1.6. Scope of study	3
1.7. Limitation	3
 2 LITERATURE REVIEW	
2.1. History of oil palm (<i>Elaeis guineensis</i>) in Malaysia	4
2.1.1. Planted area under oil palm in Malaysia	6
2.1.2. Oil palm production in Malaysia	7
2.1.3. Facto of oil palm production	7
2.1.4. Fruit	8
2.1.5. Fruit characteristic of economic importance	10
 3 RESEARCH METHODOLOGY	12
 4 RESULTS	
4.1. Plot A	22
4.2. Plot B	26
 5 DISCUSSION	30
 6 CONCLUSION AND RECOMMENDATION	35
 CITED REFERENCES	38
APPENDICES	40
CURRICULUM VITAE	92

LIST OF FIGURES

<u>Figure</u>	<u>Caption</u>	<u>Page</u>
4.1	Boxplot of bunch components in Plot A	22
4.2	Scatterplot of Plot A based on two different regression equation	25
4.3	Boxplot of bunch components in Plot B	26
4.4	Scatterplot of Plot B based on two different regression equation	29

ABSTRACT

DETERMINATION OF BUNCH COMPONENT WHICH CONTRIBUTE WITH OIL-TO-BUNCH (O/B) RATIO OF COMMERCIAL PRODUCTION IN SIME DARBY BANTING

Analysis of bunch components on oil palm (*Elaeis guineensis*) have been conducted from October 2014 to May 2015 to determine the trend of each bunch components that might contribute to the oil-to-bunch (O/B) production through all of the bunch components relationship. The raw data was obtained from Sime Darby research centre at Banting, Selangor. The raw data contains 13th year of research on the bunch analysis. The bunch analysis were collected from two different plots in order to gain more data which at the same time it will give higher amount of significant data. Although the data is taken from different plot, but they were identical characteristics such type of soil, used varieties (tenera) and also the management practices. The first analysis that have been done was trends of each bunch components as the ratio of oil-to-bunch (%) which it was measure as response variable. There are eight type of bunch components that have been considered, which are; fruit-to-bunch (F/B), mean fruit weight (MFW), mesocarp-to-fruit (M/F), kernel-to-fruit (K/F), shell-to-fruit (S/F), oil-to-dry mesocarp (O/DM), oil-to-wet mesocarp (O/WM) and kernel-to-bunch (K/B). Based on the result it can be seen the major bunch components conduct for the production was fruit-to-bunch (F/B) as it will assign other trend of bunch components allocated.